

### Problem Set #4

Sakurai and Napolitano problems:

5.20 [5.20], 5.21 [5.21]

The old (red) Sakurai (revised, 1<sup>st</sup> ed.) problems are listed in brackets.

#### 1. Helium interaction term

Show that the helium interaction term in the Hamiltonian is

$$\langle \psi_{1s}(\vec{r}_1) | \langle \psi_{1s}(\vec{r}_2) | \frac{e^2}{|\vec{r}_2 - \vec{r}_1|} | \psi_{1s}(\vec{r}_2) \rangle | \psi_{1s}(\vec{r}_1) \rangle = \frac{5}{8} Z \frac{e^2}{a_0}$$

#### 2. Bouncing quantum particle

Consider a quantum particle of mass  $m$  in a constant gravitational field  $\vec{g} = -g\hat{z}$  and constrained to move only along the  $z$ -axis, while being subject to a hard barrier at  $z=0$  from below (i.e. potential energy  $U(z \rightarrow 0) = +\infty$ ).

- Write down the Schrodinger equation for the particle and any boundary conditions that it is subject to.
- Estimate the energy of the ground state of the system using the variational method and your reasonable best choice of a trial function of your choosing.
- Estimate the energy of the first excited state.

#### 3. Minimax principle for the second excited state

Formulate the minimax principle for the second excited state and prove it.

#### 4. More variational method

$\psi_1$  and  $\psi_2$  are any two wavefunctions (not necessarily normalized and orthogonal). Consider the sub-Hilbert space spanned by these two wave functions (i.e.  $\alpha\psi_1 + \beta\psi_2$ ). The expectation of the Hamiltonian  $H$  in this subspace is given by  $\langle H \rangle = N/D$ , where

$$N = (\alpha^* \ \beta^*) \begin{pmatrix} H_{11} & H_{12} \\ H_{21} & H_{22} \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \end{pmatrix}$$

$$D = (\alpha^* \ \beta^*) \begin{pmatrix} \langle \psi_1 | \psi_1 \rangle & \langle \psi_1 | \psi_2 \rangle \\ \langle \psi_2 | \psi_1 \rangle & \langle \psi_2 | \psi_2 \rangle \end{pmatrix} \begin{pmatrix} \alpha \\ \beta \end{pmatrix}$$

$$H_{ij} = \langle \psi_i | H | \psi_j \rangle$$

Show that when  $\alpha$  and  $\beta$  are varied, the extrema of  $\langle H \rangle$  is given by the solutions of  $\lambda$  of the equation

$$\det \left[ \begin{pmatrix} H_{11} & H_{12} \\ H_{21} & H_{22} \end{pmatrix} - \lambda \begin{pmatrix} \langle \psi_1 | \psi_1 \rangle & \langle \psi_1 | \psi_2 \rangle \\ \langle \psi_2 | \psi_1 \rangle & \langle \psi_2 | \psi_2 \rangle \end{pmatrix} \right] = 0$$

These are the energy values in the sub-Hilbert space under consideration.