

Today's Topics

Monday, March 31, 2025 (Week 9, lecture 23) – Chapter 24.

A. Special Relativity review.

B. General Relativity.

C. Gravitational redshift.

D. Gravitational Waves.

E. Black holes.

Problem Set #7 is due on ExpertTA on Friday, April 4, 2025, by 9:00 AM

Midterm #2 will be on Monday, April 7, 2025.

Today's Topics

Monday, March 31, 2025 (Week 9, lecture 23) – Chapter 24.

A. Special Relativity review.

What happens when you travel close to the speed of light “ c ”

B. General Relativity.

C. Gravitational redshift.

D. Gravitational Waves.

E. Black holes.

Today's Topics

Monday, March 31, 2025 (Week 9, lecture 23) – Chapter 24.

A. Special Relativity review.

What happens when you travel close to the speed of light “ c ”

B. General Relativity.

What happens when you have very strong gravity

C. Gravitational redshift.

D. Gravitational Waves.

E. Black holes.

Special Relativity (REVIEW)

Principle of Relativity

The laws of physics are the same in all inertial reference frames.

Corollary #1

You cannot tell if you are moving (based on local measurements) in an inertial frame.

Corollary #2: Universal speed of light

The speed of light in vacuum is the same in all inertial frames, regardless of the motion of the source.

Special Relativity (REVIEW)

Principle of Relativity

The laws of physics are the same in all inertial reference frames.

Corollary #1

You cannot tell if you are moving (based on local measurements) in an inertial frame.

Corollary #2: Universal speed of light

The speed of light in vacuum is the same in all inertial frames, regardless of the motion of the source.



Length contraction & time dilation

Special Relativity (REVIEW)

Principle of Relativity

The laws of physics are the same in all inertial reference frames

Corollary #1

You cannot tell if you are moving (based on local measurements) in an inertial frame.

Corollary #2: Universal speed of light

The speed of light in vacuum is the same in all inertial frames, regardless of the motion of the source.



Length contraction & time dilation

General Relativity

Equivalence Principle

A coordinate system that is falling freely in a gravitational field is (equivalent to) an inertial frame.


Corollary

You cannot tell if you are at rest in a non-gravitational field (i.e. in a standard inertial frame) or freely falling under gravity based on local measurements.

Equivalence Principle on ISS




Curved Space-Time: light rays in 2D




Curved Space-Time

Eddington's measurement of deflection of light


- Arthur Eddington measures the deflection of starlight by the Sun.
- 1919 solar eclipse: West Africa & Brazil.
- The star appears shifted: Measurements show deflection that agrees with General Relativity.



Curved Space-Time



Curved Space-Time



[Wikipedia: Lucas Vieira Barbosa]

Gravitational Time Dilation: small heights


Clocks in a gravitational field run slower than clocks in free space.

For small changes in height Δh :

$$\frac{\Delta f}{f} = \frac{g\Delta h}{c^2}$$

f = frequency of clock

g = acceleration of gravity
= 9.8 m/s² at Earth's surface



Gravitational Time Dilation: large distances


Clocks in a gravitational field run slower than clocks in free space.

$$\frac{f_2}{f_1} = \sqrt{\frac{1 - \frac{R_s}{R_2}}{1 - \frac{R_s}{R_1}}}$$

$f_{1,2}$ = frequencies at R_1 and R_2 .

R_s = Schwarzschild radius
$$= \frac{2GM}{c^2}$$

M = mass of Earth, star, etc.




Earth

Gravitational Redshift:

Light shifts to the **red** when it escapes gravity

As light leaves the gravitational pull of Earth/star/blackhole, it loses “kinetic energy” and shifts to the red ($E_{photon} = hf$).




[<https://sites.google.com/site/salamcosmology/research/relativistic-effects-on-lss>]

Gravitational Redshift:

Light shifts to the **red** when it escapes gravity

As light leaves the gravitational pull of Earth/star/blackhole, it loses “kinetic energy” and shifts to the red ($E_{\text{photon}} = hf$).



Planck's
Constant
 $h = 6.626 \times 10^{-34} \text{ J.S}$

[<https://sites.google.com/site/salamcosmology/research/relativistic-effects-on-lss>]

Gravitational Redshift:

Light shifts to the **red** when it escapes gravity


As light leaves the gravitational pull of Earth/star/blackhole, it loses “kinetic energy” and shifts to the red ($E_{\text{photon}} = hf$).

$$\frac{\lambda_2}{\lambda_1} = \sqrt{\frac{1 - \frac{R_s}{R_2}}{1 - \frac{R_s}{R_1}}}$$

$\lambda_{1,2}$ = wavelengths at R_1 and R_2 .

$$R_s = \text{Schwarzschild radius} \\ = \frac{2GM}{c^2}$$

M = mass of Earth, star, etc.



Planck's
Constant
 $h = 6.626 \times 10^{-34} \text{ J}\cdot\text{s}$

[<https://sites.google.com/site/salamcosmology/research/relativistic-effects-on-lss>]


Earth

Gravitational Waves

- Accelerating and **orbiting** masses will emit gravitational waves.
- Gravitational waves are a consequence of the **finite speed of gravity** (*speed of light*).
 - a change in gravity's strength propagates at the speed of light.
(i.e. it's not instantaneous.)
- Only large masses emit significant gravitational waves.
 - Orbiting **black holes** and **neutron stars**.
 - Masses must be close together (i.e. fast moving) for significant emission.


Gravitational Waves

- Accelerating and **orbiting** masses will emit gravitational waves.
- Gravitational waves are a consequence of the **finite speed of gravity** (*speed of light*).
 - a change in gravity's strength propagates at the speed of light.
(i.e. it's not instantaneous.)
- Only large masses emit significant gravitational waves.
 - Orbiting **black holes** and **neutron stars**.
 - Masses must be close together (i.e. fast moving) for significant emission.
- A passing gravitational wave applies weak pulling & stretching forces along two perpendicular axes.



“+” polarization


or



“x” polarization


Gravitational Waves

- Accelerating and **orbiting** masses will emit gravitational waves.
- Gravitational waves are a consequence of the **finite speed of gravity** (*speed of light*).
 - a change in gravity's strength propagates at the speed of light.
(i.e. it's not instantaneous.)
- Only large masses emit significant gravitational waves.
 - Orbiting **black holes** and **neutron stars**.
 - Masses must be close together (i.e. fast moving) for significant emission.
- A passing gravitational wave applies weak pulling & stretching of space along two perpendicular axes (and time).



“+” polarization

or



“x” polarization

Gravitational Wave “Telescope”

LIGO: Laser Interferometer Gravitational-Wave Observatory



[<http://ligo.caltech.edu>: LIGO Livingston, LA]

Black Holes

Black hole

A celestial object whose gravity is so strong that even light cannot escape from it.

- Light emitted outside of the **event horizon** (i.e. **Schwarzschild radius**) can escape.
- Light emitted within the **event horizon** cannot escape.
- The **event horizon** / **Schwarzschild radius** defines the size and surface of a black hole.

Black Holes

Black hole

A celestial object whose gravity is so strong that even light cannot escape from it.

- Light emitted outside of the **event horizon** (i.e. **Schwarzschild radius**) can escape.
- Light emitted within the **event horizon** cannot escape.
- The **event horizon** / **Schwarzschild radius** defines the size and surface of a black hole.

$$\text{Schwarzschild radius} = R_S = \frac{2GM}{c^2}$$

*The **event horizon** is about 2-3 times smaller than the black shadow.*

Supermassive black hole at center of M87 galaxy.



[Event Horizon Telescope, www.eso.org, $\lambda=1.3$ mm]

Black Holes

Black hole

A celestial object whose gravity is so strong that even light cannot escape from it.

- Light emitted outside of the **event horizon** (i.e. **Schwarzschild radius**) can escape.
- Light emitted within the **event horizon** cannot escape.
- The **event horizon** / **Schwarzschild radius** defines the size and surface of a black hole.

$$\text{Schwarzschild radius} = R_S = \frac{2GM}{c^2}$$

*The **event horizon** is about 2-3 times smaller than the black shadow.*

Supermassive black hole at center of M87 galaxy.

[Event Horizon Telescope, www.eso.org, $\lambda=1.3$ mm]

Computer simulation of accretion disc around a black hole.

[NASA's Goddard Space Flight Center/Jeremy Schnittman]

PollEv Quiz: PollEv.com/sethaubin

2020 Nobel Prize in Physics

Black Hole Physics & Astronomy



Roger Penrose
(U. of Oxford)



Reinhard Genzel
(Max Planck Inst.)



Andrea Ghez
(UC Los Angeles)

2020 Nobel Prize in Physics

Black Hole Physics & Astronomy



Roger Penrose
(U. of Oxford)

Black hole
physics & mathematics



Reinhard Genzel
(Max Planck Inst.)




Andrea Ghez
(UC Los Angeles)

Discovery of the black hole at the center
of our Milky Way galaxy


Black Hole at center of Milky Way

The Sagittarius A* supermassive black hole



Black Hole at center of Milky Way

The Sagittarius A* supermassive black hole



[EHT Collaboration, 2022]

Mass = 4 million M_{Sun}

What happens if you fall into a Black Hole?

Stellar mass black hole

- The **Roche limit** is well outside of the event horizon.
- Any object falling towards the event horizon is **pulled apart** (spaghettified) by the strong **gravity gradient** (tidal force) of the black hole.

What happens if you fall into a Black Hole?

Stellar mass black hole

- The **Roche limit** is well outside of the event horizon.
- Any object falling towards the event horizon is **pulled apart** (spaghettified) by the strong **gravity gradient** (tidal force) of the black hole.

Supermassive black hole

- The **Roche limit** is well inside of the event horizon.
- Only after passing the event horizon is an object **pulled apart** by the black hole.

What happens if you fall into a Black Hole?

Stellar mass black hole

- The **Roche limit** is well outside of the event horizon.
- Any object falling towards the event horizon is **pulled apart** (spaghettified) by the strong **gravity gradient** (tidal force) of the black hole.

Supermassive black hole

- The **Roche limit** is well inside of the event horizon.
- Only after passing the event horizon is an object **pulled apart** by the black hole.

What happens if you watch an object fall into a black hole ?

Gravitational redshift: As the object falls its light becomes redder and eventually shifts into radio-waves.

What happens if you fall into a Black Hole?

Stellar mass black hole

- The **Roche limit** is well outside of the event horizon.
- Any object falling towards the event horizon is **pulled apart** (spaghettified) by the strong **gravity gradient** (tidal force) of the black hole.

Supermassive black hole

- The **Roche limit** is well inside of the event horizon.
- Only after passing the event horizon is an object **pulled apart** by the black hole.

What happens if you watch an object fall into a black hole ?

Gravitational redshift: As the object falls its light becomes redder and eventually shifts into radio-waves.

Gravitational time dilation: The object appears to slow down as it gets closer and closer to the event horizon.

What happens if you fall into a Black Hole?

Stellar mass black hole

- The **Roche limit** is well outside of the event horizon.
- Any object falling towards the event horizon is **pulled apart** (spaghettified) by the strong **gravity gradient** (tidal force) of the black hole.

Supermassive black hole

- The **Roche limit** is well inside of the event horizon.
- Only after passing the event horizon is an object **pulled apart** by the black hole.

What happens if you watch an object fall into a black hole ?

Gravitational redshift: As the object falls its light becomes redder and eventually shifts into radio-waves.

Gravitational time dilation: The object appears to slow down as it gets closer and closer to the event horizon.

→ Very close to the event horizon, the object becomes too redshifted to be well seen and also appears to come to a standstill.

(note: in frame of object, the object falls into black hole.)